

L SHELL XANES FOR SOLID METALS:
Ti, V, Cr, Fe, Ni, Cu

N.K. Del Grande

January 1990

Lawrence
Livermore
National
Laboratory

This is an informal report intended primarily for internal or limited external distribution. The opinions and conclusions stated are those of the author and may or may not be those of the Laboratory.

Work performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

CIRCULATION COPY
SUBJECT TO RECALL
IN TWO WEEKS

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This report has been reproduced
directly from the best available copy.

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information
P.O. Box 62, Oak Ridge, TN 37831
Prices available from (615) 576-8401, FTS 626-8401

Available to the public from the
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Rd.,
Springfield, VA 22161

L SHELL XANES FOR SOLID METALS: Ti, V, Cr, Fe, Ni, Cu

N.K. Del Grande

University of California, Lawrence Livermore National Laboratory
P.O. Box 808 (L-379) Livermore, CA 94550

Introduction

The L shell x-ray absorption near edge structure (XANES) for the 3d transition metals Ca-Cu is dominated by atomic-like behavior which is poorly understood.¹ It has been studied by electron energy loss spectroscopy (EELS) and bremsstrahlung isochromat spectra (BIS).^{2,3} More and better photoabsorption cross-section measurements are needed over the sparsely studied 400-1500 eV region to test theoretical methodologies which fuse atomic and condensed matter theories. Progress has been made reporting absolute photoabsorption cross sections taken with the spherical grating monochromator on beamline 8-2 at SSRL in 1989. These data were used in conjunction with data taken on beamline 3-4 at SSRL and beamline U14A at NSLS.^{4,7}

Experiment

Transmission data for well characterized carbon-only and carbon-metal multilayer targets (100-4000 nm thick) were recorded with a resolution of 1 eV or better for Ti, V, Cr, Fe, Ni and Cu. Absolute cross sections in Figure 1 had overall uncertainties of 10% or less. Comparisons are made with previous results.^{8,9} Away from edges, targets had photon penetration depths 1/3 to 3 mean free paths thick. Thinner targets (by a factor of 3 or 4) were used to resolve XANES shown in Figure 2.

Corrections for oxide impurities were possible since synchrotron radiation sources allowed continuous coverage across the oxygen K absorption edge at 543 eV. The determination of the target metallic areal densities (mass per unit area) was known to better than $\pm 10\%$. Targets were characterized for their total as well as component element purity, weight-per-unit area, thickness and composition, using very accurate balances, ion back-scattering, and particle-induced x-ray emission.




Figure 1. Photoabsorption cross section measurements in cm^2/g with 10% overall uncertainties are compared with the 1981 LLNL sub-keV theoretical data base⁸ and other data⁹ for five 3d solid metals: Ti, Cr, Fe, Ni, Cu. See reference [7].

Discussion

The Figure 1 comparisons of experiment with single particle Hartree Slater (HS) theory⁸ shown as a long dashed line agree for most 3d metals below and at twice the energy of the ionization threshold. The 2p-3d autoionizing resonances of Figure 2 are associated with bound-bound transitions which lead to non-radiative Auger decay. What is measured is a combination of the bound-bound photoabsorption and bound-free photoionization cross sections. The HS calculation [in 8] does not include the bound-bound component for the atom or band theory to address the behavior of the solid.

There is good qualitative agreement between L shell XANES and calculations which use a multi-configuration Dirac Fock (MCDF) model with the assumption of a statistical distribution for the initial atomic states.




Figure 2. Measured XANES for the transition metals Ti, V, Cr, Fe, Ni and Cu showing the atomic-like 2p-3d autoionizing resonances which dominate the L_3 and L_2 edge structure.⁷

Calculations were performed in intermediate coupling with configuration interactions.⁷ It is not yet clear why the statistical distribution rather than the ground state distribution for the initial states produces the measured XANES for solid metals shown in Figure 3.




Figure 3. A comparison of measurement with calculation for L shell autoionizing resonances of solid metals dominated by atomic-like behavior. (Calculation by Mau H. Chen.)

REFERENCES

1. Davis, L. C. (1986), *J. Appl. Phys.* **59**, R25-R63.
2. Fink, J., Muller-Heinzerling, M. Th., Scheerer, B., Speier, W., Hillebrecht, F.U., Fuggle, J.C., Zaanen, J. and Sawatzky, G.A., (1985), *Phys. Rev. B.*, **32** pp. 4899-4904.
3. Zaanen, J., Sawatzky, G.A., Fink, J., Speier, W., Fuggle, J.C. (1985), *Phys. Rev. B.*, **32**, pp. 4905-4913.
4. Del Grande, N.K., Tirsell, K.G., Schneider, M.B., Garrett, R.F., Kneedler, E.M., and Manson, S.T. (1987), *J. Phys. (Paris)* **48**, C9, pp.951-954.
5. Del Grande, N.K., and Tirsell, K.G. (1988), *SPIE Vol. 911 X-ray and VUV Interaction Data Bases, Calculations, and Measurements*, pp. 6-10.
6. Tirsell, K.G., and Del Grande, N.K. (1988), *SPIE Vol 911, X-ray and VUV Interaction Data Bases, Calculations, and Measurements*, pp. 146-156.
7. Del Grande, Nancy Kerr (1990), *Physica Scripta*, in press.
8. Plechaty, E., Cullen, D. and Howerton, R., (1981), Lawrence Livermore Laboratory Report, UCRL-50400, Rev. 3.
9. Del Grande, Nancy Kerr (1986), *SPIE Vol. 691, X-ray Imaging II*, pp. 2-10.

ACKNOWLEDGEMENTS

The author is indebted to K. G. Tirsell for detector improvements, to Mau H. Chen for calculations, to T. Barbee for beamline 3-4 SSRL measurements, and to P. Pianetta and his staff at SSRL for their support.